

AD \_\_\_\_\_

Award Number: W81XWH-10-1-0765

TITLE: T-Cell Immunotherapies for Treating Breast Cancer

PRINCIPAL INVESTIGATOR: Laurence Cooper, M.D., Ph.D.

CONTRACTING ORGANIZATION: The University of Texas MD Anderson Cancer Center  
Houston, TX 77030-4009

REPORT DATE: September 2011

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

<b>REPORT DOCUMENTATION PAGE</b>				<i>Form Approved</i> <b>OMB No. 0704-0188</b>	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b></small>					
<b>1. REPORT DATE</b> September 2011		<b>2. REPORT TYPE</b> Final		<b>3. DATES COVERED</b> 1 September 2010 – 31 August 2011	
<b>4. TITLE AND SUBTITLE</b>  T-Cell Immunotherapies for Treating Breast Cancer				<b>5a. CONTRACT NUMBER</b>	
				<b>5b. GRANT NUMBER</b> W81XWH-10-1-0765	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>  Laurence Cooper, M.D., Ph.D.  <b>E-Mail:</b> tforeman@mdanderson.org				<b>5d. PROJECT NUMBER</b>	
				<b>5e. TASK NUMBER</b>	
				<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  The University of Texas MD Anderson Cancer Center Houston, TX 77030-4009				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited					
<b>13. SUPPLEMENTARY NOTES</b>					
<b>14. ABSTRACT</b>  None provided.					
<b>15. SUBJECT TERMS</b> None provided.					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U			USAMRMC
			UU	19	<b>19b. TELEPHONE NUMBER</b> (include area code)

## Table of Contents

	<u>Page</u>
<b>Introduction.....</b>	<b>1</b>
<b>Body.....</b>	<b>2</b>
<b>Key Research Accomplishments.....</b>	<b>3</b>
<b>Reportable Outcomes.....</b>	<b>7</b>
<b>Conclusion.....</b>	<b>8</b>
<b>References.....</b>	<b>9</b>
<b>Appendices.....</b>	<b>16</b>

## **Introduction**

### **Breast cancer**

Breast cancer is the most commonly diagnosed cancer among U.S. women (approximately 28% or more than 1 in 4) with approximately 1 in 8 women expected to develop this malignancy over the course of her lifetime. In 2010, there are an estimated 207,000 newly diagnosed cases of invasive breast cancer resulting in about 39,800 deaths in the United States, a mortality rate higher than all other malignancies, except lung cancer. The 5-year survival rate hovers around 75-90 % for women diagnosed in the early stages with a dismal 15% for late stage breast cancer. While there are now more than 2.5 million breast cancer survivors in the U.S., in part due to better diagnostic and therapeutic efficacies, there is still plenty of room for improvement to reduce the mortality rates. Clearly, this is an important disease for women's health and significant resources are focused on developing novel and more effective treatment options.

### **Adoptive T-Cell Immunotherapy**

We proposed supplementing current breast cancer treatment options with T-cell immunotherapy which can treat malignancies resistant to conventional therapies such as surgery, chemotherapy, and radiation. As breast carcinomas grow despite a functioning immune system they develop strategies to avoid immune recognition and elimination by T cells. Gene reprogramming bestow T cells with tumor targeting capabilities and early-phase clinical trials are currently testing efficacies of T cells with redirected specificity. In addition to redirecting specificity through the expression of a chimeric antigen receptor (CAR) against c-Met, which is highly expressed in aggressive breast cancer cells, we have engineered T cells to exploit tumor hypoxia as a conditional stimulus for activation and proliferation. By programming T cells to activate only in pathological tumor hypoxia (1% O<sub>2</sub>), our approach seeks to avoid deleterious off-target effects, bypassing physiological normoxic (20% O<sub>2</sub>) and normal c-Met<sup>+</sup> tissues.

### **Hypoxic tumor microenvironment and c-Met**

We selected c-Met, a hypoxia-inducible antigen widely expressed in breast cancer, as the target for redirected T-cell specificity for the following reasons:

- 1) Poor prognosis of c-Met<sup>+</sup> breast tumors
- 2) Widely expressed in chemo-resistant and hypoxic breast tumors

### 3) c-Met signaling pathway significantly affects breast tumor survival

Our approach is to exploit the hypoxic breast tumor microenvironment to devise a robust tumor specific T-cell immunotherapy. Relapsed breast tumors are hypoxic and resistant to conventional chemotherapy and radiotherapy regimens. These hypoxic tumors are prone to metastasis leading to widespread treatment failure. Hypoxic breast tumors express high levels of c-Met, a hypoxia-inducible tyrosine kinase receptor for hepatocyte growth factor/scatter factor (HGF/SF), and a multipotent cytokine that induces invasive and metastatic growth. c-Met activation triggers mitogenic, morphogenic, and angiogenic changes in breast tumors, invariably associated with extremely poor prognosis, reduced survival, and high metastatic risks.

## Body

We hypothesize that a combination of immunotherapy which exploits the tumor microenvironment would probably be a innovative and novel alternative to supplement current treatment modalities

## Goals

To develop chimeric antigen receptors (CARs) to target antigens on breast cancer cells. Our approach focuses on exploiting the hypoxic tumor microenvironment to conditionally activate reprogrammed T cells specific for the c-Met antigen. Specifically, we want to (1) develop a CAR specific for c-Met antigen to reprogram T cells for antigen-dependent killing of breast cancer cells, and (2) develop a CAR to target breast cancer cells that can be conditionally expressed under hypoxia.

## CAR constructs

c-Met specific CAR was constructed by fusing the single-chain variable fragments (scFv) of a mouse anti-human c-Met antibody to the human IgG4 Fc domain and signaling endodomains of human CD28 and CD3 (CD247). In addition, to render the CAR oxygen-sensitive, an oxygen-dependent degradation domain from Hypoxia Inducible Factor-1 (HIF-1, the master transcriptional regulator of cellular hypoxia response) was incorporated into the c-Met CAR yielding c-Met-ODDD CAR, which is oxygen-sensitive. Both the c-Met CAR and c-Met-ODDD CAR were cloned into *Sleeping Beauty* (SB) transposition vector (transposon DNA, schematic in Appendix). The expression cassettes were flanked by inverted

repeats/direct repeats (IR/DR) for specific recognition by SB transposase, which cuts and pastes the transposon DNA into genomic T cell DNA.

### **Growing of CAR on aAPC platform**

Peripheral blood mononuclear cells (PBMC) were isolated from normal donors by venipuncture and Ficoll fractionation. These cells are nucleofected using Amaxa T cell transfection with SB transposase and transposon constructs. The cells were then stimulated using  $\gamma$ -irradiated artificial antigen presenting cells (aAPC), a K562-based HLA<sup>-ve</sup> cell line constitutively expressing c-Met antigen, CD86, CD137L, IL-15, and CD64. These co-stimulatory molecules ensure optimal T cell expansion.

### **Testing of CAR expression based on oxygen condition**

T cells were expanded on 1% (hypoxic condition) or 20% (normoxic condition) oxygen supplemented with 5% CO<sub>2</sub> in a humidified environment at 37°C. CAR expression were determined by Fluorescent-Activated Cell Sorting (FACS). Cell numbers were enumerated and fold expansion calculated over a 4-week (4 x 7 days stimulations) period.

### **Testing of CAR on breast cancer cell lines**

c-Met expression was verified on breast cancer cell lines and co-cultures were performed to assay for CAR-directed T cells killing of target tumor cells under either low or normal oxygen conditions (1%, or 20% O<sub>2</sub>).


## **Key Research Accomplishments**

We capitalized on the triple associations of hypoxia, c-Met, and breast carcinoma to develop novel and specific immunotherapy for enhanced therapeutic efficacy. Successful targeting of breast cancer will involve not only redirected specificity, but also modifying T cells to overcome the hostile/hypoxic tumor microenvironment:


Research accomplishments include:

- (i) We can use the oxygen-dependent degradation domain (ODDD) to render the expression of red-fluorescent protein sensitive to hypoxia (Figure 1).

- (ii) We can fuse ODDD to c-Met-specific CAR to render T cells capable of only expressing the CAR-ODDD under conditions of hypoxia, but not normoxia (Figure 1). Control c-Met-specific CAR is expressed under hypoxia and normoxia.
- (iii) We can propagate c-Met-specific CAR-ODDD to large and clinically-sufficient numbers under hypoxia on  $\gamma$ -irradiated c-Met<sup>+</sup> aAPC (Figure 2).
- (iv) We can demonstrate re-directed effector function of c-Met-ODDD<sup>+</sup> T cells based on selected killing of hypoxic tumor cells expressing c-Met (Figure 3).




**Figure 1:** (A) Expression (in K562 cells) of the red-fluorescent protein AsRed-2, fused to ODDD, under hypoxia (1%  $O_2$ ) and then loss of expression under normoxia (20%  $O_2$ ), and followed by re-expression of AsRed2-ODDD as the oxygen tension was reduced. (In these preliminary data, not all the cells express AsRed2-ODDD.) Yellow circles highlight cell fluorescence indicating molecular sensing of  $O_2$  and AsRed-2 expression. (B) Expression of c-Met-specific CAR (without fusion to ODDD) under normoxia and hypoxia (top). The human HIF-1 ODDD was fused in frame to the 3' end of the c-Met-specific CAR (CAR-



**Figure 2:** Selective propagation of c-Met-specific CAR-ODDD<sup>+</sup> T cells can be numerically expanded on c-Met<sup>+</sup> aAPC under hypoxia (1%  $O_2$ ).





**Figure 3:** Chromium release assay of c-Met-specific CAR-ODDD<sup>+</sup> T cells under conditions of normoxia and hypoxia. The tumor cells (K562) were transfected to enforce expression of c-met on the cell surface independent of O<sub>2</sub> concentration. The T cells conditionally express the CAR-ODDD under conditions of hypoxia.

Reportable Outcomes


The ability to genetically manipulate T cells and expand them to clinically relevant numbers *ex vivo* provides new opportunities to enhance their biologic activity in vivo.


Our data demonstrate that:

- (i) c-Met CAR<sup>+</sup> T cells are able to kill c-Met<sup>+</sup> breast cancer cells only (Figure 4).
- (ii) c-Met ODDD CAR<sup>+</sup> T cells are able to kill c-Met<sup>+</sup> breast cancer cells only in low oxygen conditions, such as tumor environments (Figure 5).

Table 1. c-Met expression in breast cancer cells

Breast cancer cell lines:	c-Met expression:
T47D	low / negative
MCF-7	low/ negative
MDA-MB231	positive
SkBr-3	positive





## Conclusions

We have generated a CAR coupled to a molecular sensor for oxygen that is capable of targeting c-Met which can be useful in the treatment of breast cancer. Conditional activation of T cells can be a useful way of avoiding unwanted effects with antigens which have a widespread expression profile. Oxygen-sensitive CARs can be used to activate T cells only in tumor microenvironment leaving normal c-Met<sup>+</sup> epithelial cells untouched.

c-Met overexpressions are also detected in other malignancies so our platform technology can also be useful in tumors with pathologic HGF/c-Met signaling axis. In addition, this technology can be adapted to target specific antigens of choice for other tumor types.

To our knowledge, no publication is found detailing therapeutic development conditionally targeting c-Met in breast carcinomas. We are therefore at the cutting edge forefront in our attempt to leverage our platform technology to conditionally reprogram T cells to target c-Met in breast cancer.

## References/Bibliography of Publications and Meeting Abstracts

Bertotti A, Comoglio PM, Trusolino L. Beta4 integrin is a transforming molecule that unleashes Met tyrosine kinase tumorigenesis. *Cancer Res.* 65, 10674-9 (2005)

Beviglia, L., Matsumoto, K., Lin, C. S., Ziober, B. L. & Kramer, R. H. Expression of the c-Met/HGF receptor in human breast carcinoma: correlation with tumor progression. *Int J Cancer* 74, 301-9. (1997).

Bigelow RL, Cardelli JA. The green tea catechins, (-)-Epigallocatechin-3-gallate (EGCG) and (-)-Epicatechin-3-gallate (ECG), inhibit HGF/Met signaling in immortalized and tumorigenic breast epithelial cells. *Oncogene.* 25, 1922-30 (2006)

Camp, R. L., Rimm, E. B. & Rimm, D. L. Met expression is associated with poor outcome in patients with axillary lymph node negative breast carcinoma. *Cancer* 86, 2259-65. (1999).

Carracedo A, Egervari K, Salido M, Rojo F, Corominas JM, Arumi M, Corzo C, Tusquets I, Espinet B, Rovira A, Albanell J, Szollosi Z, Serrano S, Solé F. FISH and immunohistochemical status of the hepatocyte growth factor receptor (c-Met) in 184 invasive breast tumors. *Breast Cancer Res.* 11, 402 (2009)

Castro NE, Lange CA. Breast tumor kinase and extracellular signal-regulated kinase 5 mediate Met receptor signaling to cell migration in breast cancer cells. *Breast Cancer Res.* 12, R60 (2010)

Charafe-Jauffret E, Ginestier C, Monville F, Finetti P, Adelaide J, Cervera N, Fekairi S, Xerri L, Jacquemier J, Birnbaum D, Bertucci F. Gene expression profiling of breast cell lines identifies potential new basal markers. *Oncogene.* 25, 2273-84 (2006)

Chen CC, Hou MF, Wang JY, Chang TW, Lai DY, Chen YF, Hung SY, Lin SR. Simultaneous detection of multiple mRNA markers CK19, CEA, c-Met, Her2/neu and hMAM with membrane array, an innovative technique with a great potential for breast cancer diagnosis. *Cancer Lett.* 2005 Nov 11; [Epub ahead of print]

Chen HH, Su WC, Lin PW, Guo HR, Lee WY. Hypoxia-inducible factor-1alpha correlates with MET and metastasis in node-negative breast cancer. *Breast Cancer Res Treat.* 103, 167-75 (2007)

Cheng N, Chytil A, Shyr Y, Joly A, Moses HL. Enhanced hepatocyte growth factor signaling by type II transforming growth factor-beta receptor knockout fibroblasts promotes mammary tumorigenesis. *Cancer Res.* 67, 4869-77 (2007)

Coskun U, Bukan N, Sancak B, Gunel N, Ozenirler S, Unal A, Yucel A. Serum hepatocyte growth factor and interleukin-6 levels can distinguish patients with primary or metastatic liver tumors from those with benign liver lesions. *Neoplasma* 51, 209-13 (2004)

Edakuni, G., Sasatomi, E., Satoh, T., Tokunaga, O. & Miyazaki, K. Expression of the hepatocyte growth factor/c-Met pathway is increased at the cancer front in breast carcinoma. *Pathol Int* 51, 172-8. (2001).

Eichbaum MH, de Rossi TM, Kaul S, Bruckner T, Schneeweiss A, Sohn C. Serum levels of hepatocyte growth factor/scatter factor in patients with liver metastases from breast cancer. *Tumour Biol.* 28, 36-44 (2007)

Filigheddu N, Cutrupi S, Porporato PE, Riboni F, Baldanzi G, Chianale F, Fortina E, Piantanida P, De Bortoli M, Vacca G, Graziani A, Surico N. Diacylglycerol kinase is required for HGF-induced invasiveness and anchorage-independent growth of MDA-MB-231 breast cancer cells. *Anticancer Res.* 27, 1489-92 (2007)

Finkbeiner MR, Astanehe A, To K, Fotovati A, Davies AH, Zhao Y, Jiang H, Stratford AL, Shadeo A, Boccaccio C, Comoglio P, Mertens PR, Eirew P, Raouf A, Eaves CJ, Dunn SE. Profiling YB-1 target genes uncovers a new mechanism for MET receptor regulation in normal and malignant human mammary cells. *Oncogene.* 28, 1421-31 (2009)

Furlan A, Vercamer C, Desbiens X, Pourtier A. Ets-1 triggers and orchestrates the malignant phenotype of mammary cancer cells within their matrix environment. *J Cell Physiol.* 215, 782-93 (2008)

Garcia S, Dales JP, Charafe-Jauffret E, Carpentier-Meunier S, Andrac-Meyer L, Jacquemier J, Andonian C, Lavaut MN, Allasia C, Bonnier P, Charpin C. Overexpression of c-Met and of the transducers PI3K, FAK and JAK in breast carcinomas correlates with shorter survival and neoangiogenesis. *Int J Oncol.* 31, 49-58 (2007)

Garcia S, Dalès JP, Charafe-Jauffret E, Carpentier-Meunier S, Andrac-Meyer L, Jacquemier J, Andonian C, Lavaut MN, Allasia C, Bonnier P, Charpin C. Poor prognosis in breast carcinomas correlates with increased expression of targetable CD146 and c-Met and with proteomic basal-like phenotype. *Hum Pathol.* 38, 830-41 (2007)

Garcia S, Dalès JP, Jacquemier J, Charafe-Jauffret E, Birnbaum D, Andrac-Meyer L, Lavaut MN, Allasia C, Carpentier-Meunier S, Bonnier P, Charpin-Taranger C. c-Met overexpression in inflammatory breast carcinomas: automated quantification on tissue microarrays. *Br J Cancer.* 96, 329-35 (2007)

Götte M, Kersting C, Radke I, Kiesel L, Wülfing P. An expression signature of syndecan-1 (CD138), E-cadherin and c-met is associated with factors of angiogenesis and lymphangiogenesis in ductal breast carcinoma in situ. *Breast Cancer Res.* 9, R8 (2007)

Götte M, Kersting C, Radke I, Kiesel L, Wülfing P. An expression signature of syndecan-1 (CD138), E-cadherin and c-met is associated with factors of angiogenesis and lymphangiogenesis in ductal breast carcinoma in situ. *Breast Cancer Res.* 9, R8 (2007)

Graveel CR, DeGroot JD, Su Y, Koeman J, Dykema K, Leung S, Snider J, Davies SR, Swiatek PJ, Cottingham S, Watson MA, Ellis MJ, Sigler RE, Furge KA, Vande Woude GF. Met induces diverse mammary carcinomas in mice and is associated with human basal breast cancer. *Proc Natl Acad Sci U S A.* 106, 12909-14 (2009)

Greenberg R, Barnea Y, Schneebaum S, Kashtan H, Kaplan O, Skornik Y. Detection of hepatocyte growth factor/scatter factor receptor (c-Met) and MUC1 from the axillary fluid drainage in patients after breast cancer surgery. *Isr Med Assoc J* 5:649-52 (2003).

Greenberg, R, Schwartz, I, Skornick Y, & Kaplan O. Detection of hepatocyte growth factor/scatter factor receptor (c-Met) in axillary drainage after operations for breast cancer using reverse transcriptase-polymerase chain reaction. *Breast Cancer Res* 5:71-76 (2003).

Hiscox S, Jordan NJ, Jiang W, Harper M, McClelland R, Smith C, Nicholson RI. Chronic exposure to fulvestrant promotes overexpression of the c-Met receptor in breast cancer cells: implications for tumour-stroma interactions. *Endocr Relat Cancer.* 13, 1085-99 (2006)

Jedezsko C, Victor BC, Podgorski I, Sloane BF. Fibroblast hepatocyte growth factor promotes invasion of human mammary ductal carcinoma in situ. *Cancer Res.* 69, 9148-55 (2009)

Jin, L. et al. Expression of scatter factor and c-met receptor in benign and malignant breast tissue. *Cancer* 79, 749-60. (1997).

Jones BA, Kasl SV, Howe CL, Lachman M, Dubrow R, Curnen MM, Soler-Vila H, Beeghly A, Duan F, Owens P. African-American/White differences in breast carcinoma: p53 alterations and other tumor characteristics. *Cancer* 101, 1293-301 (2004)

Kang, J. Y. et al. Tissue microarray analysis of hepatocyte growth factor/Met pathway components reveals a role for Met, matriptase, and hepatocyte growth factor activator inhibitor 1 in the progression of node-negative breast cancer. *Cancer Res* 63, 1101-5 (2003).

Khoury H, Naujokas MA, Zuo D, Sangwan V, Frigault MM, Petkiewicz S, Dankort DL, Muller WJ, Park M. HGF converts ErbB2/Neu epithelial morphogenesis to cell invasion. *Mol Biol Cell*. 16 550-61(2005)

Klosek SK, Nakashiro K, Hara S, Goda H, Hasegawa H, Hamakawa H. CD151 regulates HGF-stimulated morphogenesis of human breast cancer cells. *Biochem Biophys Res Commun*. 379, 1097-100 (2009)

Lee WJ, Chen WK, Wang CJ, Lin WL, Tseng TH. Apigenin inhibits HGF-promoted invasive growth and metastasis involving blocking PI3K/Akt pathway and beta 4 integrin function in MDA-MB-231 breast cancer cells. *Toxicol Appl Pharmacol*. 226, 178-91 (2007)

Lee WY, Chen HH, Chow NH, Su WC, Lin PW, Guo HR. Prognostic significance of co-expression of RON and MET receptors in node-negative breast cancer patients. *Clin Cancer Res*. 11, 2222-8 (2005)

Lee WY, Su WC, Lin PW, Guo HR, Chang TW, Chen HH. Expression of S100A4 and Met: potential predictors for metastasis and survival in early-stage breast cancer. *Oncology* 66, 429-38 (2004)

Leibowitz-Amit R, Tsarfaty G, Abargil Y, Yerushalmi GM, Horev J, Tsarfaty I. Mimp, a mitochondrial carrier homologue, inhibits Met-HGF/SF-induced scattering and tumorigenicity by altering Met-HGF/SF signaling pathways. *Cancer Res*. 66, 8687-97 (2006)

Lengyel E, Prechtel D, Resau JH, Gauger K, Welk A, Lindemann K, Salanti G, Richter T, Knudsen B, Vande Woude GF, Harbeck N. C-Met overexpression in node-positive breast cancer identifies patients with poor clinical outcome independent of Her2/neu. *Int J Cancer* 113, 678-82 (2005)

Lindemann K, Harbeck N, Lengyel E, Resau JH. A special key for unlocking the door to targeted therapies of breast cancer. *ScientificWorldJournal*. 8, 905-8 (2008)

Lindemann K, Resau J, Nährig J, Kort E, Leeser B, Annecke K, Welk A, Schäfer J, Vande Woude GF, Lengyel E, Harbeck N. Differential expression of c-Met, its ligand HGF/SF and HER2/neu in DCIS and adjacent normal breast tissue. *Histopathology*. 51, 54-62 (2007)

Matteucci E, Locati M, Desiderio MA. Hepatocyte growth factor enhances CXCR4 expression favoring breast cancer cell invasiveness. *Exp Cell Res*. 310 176-85 (2005)

Matteucci E, Ridolfi E, Desiderio MA. Hepatocyte growth factor differently influences Met-E-cadherin phosphorylation and downstream signaling pathway in two models of breast cells. *Cell Mol Life Sci.* 63, 2016-26 (2006)

Morozov VM, Massoll NA, Vladimirova OV, Maul GG, Ishov AM. Regulation of c-met expression by transcription repressor Daxx. *Oncogene.* 27, 2177-86 (2008)

Mueller KL, Hunter LA, Ethier SP, Boerner JL. Met and c-Src cooperate to compensate for loss of epidermal growth factor receptor kinase activity in breast cancer cells. *Cancer Res.* 68, 3314-22 (2008)

Mueller KL, Hunter LA, Ethier SP, Boerner JL. Met and c-Src cooperate to compensate for loss of epidermal growth factor receptor kinase activity in breast cancer cells. *Cancer Res.* 68, 3314-22 (2008)

Nagy, J. et al. Hepatocyte growth factor/scatter factor expression and c-met in primary breast cancer. *Surg Oncol* 5, 15-21. (1996).

Parr C, Watkins G, Mansel RE, Jiang WG. The hepatocyte growth factor regulatory factors in human breast cancer. *Clin Cancer Res* 10:202-11 (2004).

Ponzo MG, Lesurf R, Petkiewicz S, O'Malley FP, Pinnaduwa D, Andrulis IL, Bull SB, Chughtai N, Zuo D, Souleimanova M, Germain D, Omeroglu A, Cardiff RD, Hallett M, Park M. Met induces mammary tumors with diverse histologies and is associated with poor outcome and human basal breast cancer. *Proc Natl Acad Sci U S A.* 106, 12903-8 (2009)

Pozner-Moulis S, Cregger M, Camp RL, Rimm DL. Antibody validation by quantitative analysis of protein expression using expression of Met in breast cancer as a model. *Lab Invest.* 87, 251-60 (2007)

Pozner-Moulis S, Pappas DJ, Rimm DL. Met, the hepatocyte growth factor receptor, localizes to the nucleus in cells at low density. *Cancer Res.* 66, 7976-82 (2006)

Previdi S, Maroni P, Matteucci E, Broggin M, Bendinelli P, Desiderio MA. Interaction between human-breast cancer metastasis and bone microenvironment through activated hepatocyte growth factor/Met and beta-catenin/Wnt pathways. *Eur J Cancer.* 46, 1679-91 (2010)

Pruefer FG, Unger-Saldaña K, Mohamier L, Ramírez M, Suárez J, Ramirez MT, Maldonado V, Meléndez-Zajgla J. Tumor HGF lacks prognostic significance in Mexican breast cancer patients. *J Exp Clin Cancer Res.* 25, 357-64 (2006)



Ridolfi E, Matteucci E, Maroni P, Desiderio MA. Inhibitory effect of HGF on invasiveness of aggressive MDA-MB231 breast carcinoma cells, and role of HDACs. *Br J Cancer*. 99, 1623-34. Epub 2008 Oct 21 (2008)

Sam MR, Elliott BE, Mueller CR. A novel activating role of SRC and STAT3 on HGF transcription in human breast cancer cells. *Mol Cancer*. 6, 69 (2007)

Shattuck DL, Miller JK, Carraway KL 3rd, Sweeney C. Met receptor contributes to trastuzumab resistance of Her2-overexpressing breast cancer cells. *Cancer Res*. 68, 1471-7 (2008)

Shattuck DL, Miller JK, Laederich M, Funes M, Petersen H, Carraway KL 3rd, Sweeney C. LRIG1 is a novel negative regulator of the Met receptor and opposes Met and Her2 synergy. *Mol Cell Biol*. 27, 1934-46 (2007)

Sheen-Chen SM, Liu YW, Eng HL, Chou FF. Serum levels of hepatocyte growth factor in patients with breast cancer. *Cancer Epidemiol Biomarkers Prev*. 14 715-7 (2005)

Singletery K, Ellington A. Genistein suppresses proliferation and MET oncogene expression and induces EGR-1 tumor suppressor expression in immortalized human breast epithelial cells. *Anticancer Res*. 26, 1039-48 (2006)

Smolen GA, Muir B, Mohapatra G, Barmettler A, Kim WJ, Rivera MN, Haserlat SM, Okimoto RA, Kwak E, Dahiya S, Garber JE, Bell DW, Sgroi DC, Chin L, Deng CX, Haber DA. Frequent met oncogene amplification in a Brca1/Trp53 mouse model of mammary tumorigenesis. *Cancer Res*. 66, 3452-5 (2006)

Syed DN, Afaq F, Sarfaraz S, Khan N, Kedlaya R, Setaluri V, Mukhtar H. Delphinidin inhibits cell proliferation and invasion via modulation of Met receptor phosphorylation. *Toxicol Appl Pharmacol*. 231, 52-60 (2008)

Takahashi K, Suzuki K. Requirement of kinesin-mediated membrane transport of WAVE2 along microtubules for lamellipodia formation promoted by hepatocyte growth factor. *Exp Cell Res*. 314, 2313-22 (2008)

Taraseviciute A, Vincent BT, Schedin P, Jones PL. Quantitative analysis of three-dimensional human mammary epithelial tissue architecture reveals a role for tenascin-C in regulating c-met function. *Am J Pathol*. 176, 827-38 (2010)

Tolgay Ocal, I, Dolled-Filhart, M, D'Aquila, TG, Camp, RL, & Rimm, DL. Tissue microarray-based studies of patients with lymph node negative breast carcinoma show that met expression is associated with worse outcome but is not correlated with epidermal growth factor family receptors. *Cancer* 97:1841-8 (2003).

Tsarfaty, I. et al. Alteration of Met protooncogene product expression and prognosis in breast carcinomas. *Anal Quant Cytol Histol* 21, 397-408. (1999).

Tuck, A. B., Park, M., Sterns, E. E., Boag, A. & Elliott, B. E. Coexpression of hepatocyte growth factor and receptor (Met) in human breast carcinoma. *Am J Pathol* 148, 225-32 (1996).

Welm AL, Kim S, Welm BE, Bishop JM. MET and MYC cooperate in mammary tumorigenesis. *Proc Natl Acad Sci U S A.* 102, 4324-9 (2005)

Wright TG, Singh VK, Li JJ, Foley JH, Miller F, Jia Z, Elliott BE. Increased production and secretion of HGF alpha-chain and an antagonistic HGF fragment in a human breast cancer progression model. *Int J Cancer.* 125, 1004-15 (2009)

Xu H, Washington S, Verderame MF, Manni A. Role of non-receptor and receptor tyrosine kinases (TKs) in the antitumor action of alpha-difluoromethylornithine (DFMO) in breast cancer cells. *Breast Cancer Res Treat.* 112, 255-61 (2008)

Yamashita, J. et al. Immunoreactive hepatocyte growth factor is a strong and independent predictor of recurrence and survival in human breast cancer. *Cancer Res* 54, 1630-3 (1994).

Yao, Y. et al. Scatter factor protein levels in human breast cancers: clinicopathological and biological correlations. *Am J Pathol* 149, 1707-17 (1996).

Yin YL, Chen HL, Kuo HM, He SP. NK3 and NK4 of HGF enhance filamin production via STAT pathway, but not NK1 and NK2 in human breast cancer cells. *Acta Pharmacol Sin.* 29, 728-35 (2008)


## List of Personnel Supported by Grant

Sonny Oon Ang

Tijuan Mi

## Appendices

Schematic of Sleeping Beauty Transposition system consisting of a transposase expression plasmid and a transposon DNA vector.



Left Panel: plasmid construct for SB11 transposase. Right Pane: DNA vector of c-Met specific chimeric antigen receptor (CAR)